arXiv:1505.05771v1 [math.CO] 21 May 2015

Cayley graphs of more than one abelian group

Edward Dobson

Department of Mathematics and Statistics
Mississippi State University
PO Drawer MA Mississippi State, MS 39762
and
IAM, University of Primorska
Muzejska trg 2
Koper 6000, Slovenia

Joy Morris[1]

Department of Mathematics and Computer Science
University of Lethbridge
Lethbridge, AB. T1K 3M4. Canada

Abstract

We show that for certain integers n, the problem of whether or not a Cayley digraph Γ of $\mathbb{Z}_n$ is also isomorphic to a Cayley digraph of some other abelian group G of order n reduces to the question of whether or not a natural subgroup of the full automorphism group contains more than one regular abelian group up to isomorphism (as opposed to the full automorphism group). A necessary and sufficient condition is then given for such circulants to be isomorphic to Cayley digraphs of more than one abelian group, and an easy-to-check necessary condition is provided.

---

1. Introduction

It is well known that a Cayley digraph of a group G may also be isomorphic to a Cayley digraph of group H where G and H are not isomorphic. A natural question is then to determine exactly when a Cayley digraph is isomorphic to a Cayley digraph of a nonisomorphic group. Perhaps the first work on this problem was by Joseph in 1995 [10] where she determined necessary and sufficient conditions for a Cayley digraph of order p^2, p a prime, to be isomorphic to a Cayley digraph of both groups of order p^2 (see [8, Lemma 4] for a group theoretic version of this result). The second author [16] subsequently extended this result and determined necessary and sufficient conditions for a Cayley digraph of $\mathbb{Z}_{p^k}$, $k \geq 1$ and p an odd prime, to be isomorphic to a Cayley digraph of some other abelian group (see Theorem 1.4 for the statement of this result). Additionally,

---

[1] The work of this author is supported in part by a research grant from NSERC.

she found necessary and sufficient conditions for a Cayley digraph of $\mathbb{Z}_{p^k}$ to be isomorphic to a Cayley digraph of any group of order p^k. The equivalent problem for $p = 2$ was solved by Kovács and Servatius [11]. Digraphs of order pq that are Cayley graphs of both groups of order pq, where $q \mid (p-1)$ and p, q are distinct primes were determined by the first author in [4, Theorem 3.4]. Finally, Marušič and the second author studied the question of which normal circulant graphs of square-free order are also Cayley graphs of a nonabelian group [14].

We show in this paper that for some values of n, we can reduce the problem of which circulant digraphs of order n are also Cayley digraphs of some other abelian group of order n, to the prime-power case previously solved by the second author. Specifically, let $n = p_1^{a_1} p_2^{a_2} \ldots p_r^{a_r}$ and let $k = p_1 \ldots p_r$, where each p_i is prime. Then the reduction works if $\gcd(k, \varphi(k)) = 1$.

In the remainder of this section, we state the second author's result (Theorem 1.4), first providing the necessary definitions for that statement. In Section 2, we provide the necessary group theoretic results to prove our main theorem. In Section 3, we provide the necessary graph theoretic results to prove our main result, which is Corollary 3.12.

Definition 1.1. *Let* G *be a group and* $S \subset G$*. Define a* **Cayley digraph of** G*, denoted* $\mathrm{Cay}(G, S)$*, to be the digraph with* $V(\mathrm{Cay}(G, S)) = G$ *and* $E(\mathrm{Cay}(G, S)) = \{(g, gs) : g \in G, s \in S\}$*. We call* S *the* **connection set** *of* $\mathrm{Cay}(G, S)$*.*

Definition 1.2. *Let* Γ_1 *and* Γ_2 *be digraphs. The* **wreath product of** Γ_1 **and** Γ_2*, denoted* $\Gamma_1 \wr \Gamma_2$*, is the digraph with vertex set* $V(\Gamma_1) \times V(\Gamma_2)$ *and edges* $(u, v)(u, v')$ *for* $u \in V(\Gamma_1)$ *and* $vv' \in E(\Gamma_2)$ *or* $(u, v)(u', v')$ *where* $uu' \in E(\Gamma_1)$ *and* $v, v' \in V(\Gamma_2)$*.*

Definition 1.3. *Let* G *be a permutation group acting on* X *and* H *a permutation group acting on* Y*. Define the* **wreath product of** G **and** H*, denoted* $G \wr H$*, to be the set of all permutations of* $X \times Y$ *of the form* $(x, y) \mapsto (g(x), h_x(y))$*.*

We caution the reader that these definitions of wreath products are not completely standard, in that some mathematicians use $H \wr G$ for what we have defined as $G \wr H$, and similarly use $\Gamma_2 \wr \Gamma_1$ for our $\Gamma_1 \wr \Gamma_2$. Both orderings appear in the literature.

We will often times consider wreath products of multiple digraphs or groups, and sometimes the specific digraphs or groups are unimportant. In this circumstance, rather than write out and define the digraphs or groups, we will just say that a graph Γ or group G is a **multiwreath product**. Formally, a digraph Γ is a multiwreath product if there exists digraphs $\Gamma_1, \ldots, \Gamma_r$ such that $\Gamma = \Gamma_1 \wr \Gamma_2 \wr \cdots \wr \Gamma_r$, and a group G is a multiwreath product if there exists groups $G_1, \ldots, G_r$ such that $G = G_1 \wr G_2 \wr \cdots \wr G_r$.

Following [16], define a partial order on the set of abelian groups of order p^n as follows: We say $G \preceq_p H$ if there is a chain $H_1 < H_2 < \ldots < H_m = H$ of subgroups of H such that $H_1, H_2/H_1, \ldots, H_m/H_{m-1}$ are all cyclic, and


$\mathbb{Z}_{p^5}$
$\mathbb{Z}_{p^4} \times \mathbb{Z}_p$
$\mathbb{Z}_{p^2} \times \mathbb{Z}_{p^3}$
$\mathbb{Z}_{p^2}^2 \times \mathbb{Z}_p$
$\mathbb{Z}_{p^3} \times \mathbb{Z}_p^2$
$\mathbb{Z}_{p^2} \times \mathbb{Z}_p^3$
$\mathbb{Z}_p^5$

Figure 1: The partial order for groups of order p^5

$$G \cong H_1 \times \frac{H_2}{H_1} \times \ldots \times \frac{H_m}{H_{m-1}}.$$

There is an equivalent definition for this partial order. We say that a string of integers $i_1, \ldots, i_m$ is a **subdivision** of the string of integers $j_1, \ldots, j_{m_0}$ if there is some permutation δ of $\{1, ..., m\}$ and some strictly increasing sequence of integers $0 = k_0, \ldots, k_{m_0} = m$ such that $i_{\delta(k_s+1)} + \ldots + i_{\delta(k_{s+1})} = j_{s+1}$, $0 \leq s \leq m_0 - 1$. Now, $G \preceq_p H$ precisely if $G \cong \mathbb{Z}_{p^{i_1}} \times \mathbb{Z}_{p^{i_2}} \times \ldots \times \mathbb{Z}_{p^{i_m}}$ and $H \cong \mathbb{Z}_{p^{j_1}} \times \mathbb{Z}_{p^{j_2}} \times \ldots \times \mathbb{Z}_{p^{j_{m_0}}}$ where $i_1, ..., i_m$ is a subdivision of $j_1, \ldots, j_{m_0}$. In Figure 1 this partial order is depicted for abelian groups of order p^5.

The following result was proven in [16] (see also [15]) in the case where p is an odd prime and in [11] when $p = 2$.

Theorem 1.4. *Let $\Gamma = \mathrm{Cay}(G, S)$ be a Cayley digraph on an abelian group G of order p^k, where p is prime. Then the following are equivalent:*

1. *The digraph Γ is isomorphic to a Cayley digraph on both $\mathbb{Z}_{p^k}$ and H, where H is an abelian group with $|H| = p^k$, say $H = \mathbb{Z}_{p^{k_1}} \times \mathbb{Z}_{p^{k_2}} \times \ldots \times \mathbb{Z}_{p^{k_{m_0}}}$, where $k_1 + \ldots + k_{m_0} = k$.*
2. *There exist a chain of subgroups $G_1 \leq \ldots \leq G_{m-1}$ in G such that*
   (a) *$G_1, G_2/G_1, \ldots, G/G_{m-1}$ are cyclic groups;*
   (b) *$G_1 \times G_2/G_1 \times \ldots \times G/G_{m-1} \preceq_p H$;*
   (c) *For all $s \in S \backslash G_i$, we have $sG_i \subseteq S$, for $i = 1, ..., m-1$. (That is, $S \backslash G_i$ is a union of cosets of G_i.)*
3. *There exist Cayley digraphs $U_1, ..., U_m$ on cyclic p-groups $H_1, ..., H_m$ such that $H_1 \times \ldots \times H_m \preceq_p H$ and $\Gamma \cong U_m \wr \ldots \wr U_1$.*
   These in turn imply:

4. Γ *is isomorphic to Cayley digraphs on every abelian group of order* p^k *that is greater than* H *in the partial order.*

2. Group Theoretic Results

For any terms from permutation group theory that are not defined in this paper, see [2]. In the section we collect all permutation group theoretic results that we will need for our main result.

Theorem 2.1 (Theorem 3 of [9], or Corollary 1.2 of [12]). *A primitive permutation group* K *acting on* Ω *of finite degree* n *has a cyclic regular subgroup if and only if one of the following holds:*

1. $\mathbb{Z}_p \leq K \leq \mathrm{AGL}(1,p)$, *where* $n = p$ *is prime;*
2. $K = \mathrm{Sym}(n)$ *for some* n, *or* $K = \mathrm{Alt}(n)$ *for some odd* n*;*
3. $\mathrm{PGL}(d,q) \leq K \leq \mathrm{P\Gamma L}(d,q)$ *where* $n = (q^d-1)/(q-1)$ *for some* $d \geq 2$*;*
4. $K = \mathrm{PSL}(2,11)$, M_{11} *or* M_{23} *where* $n = 11$, 11 *or* 23 *respectively.*

Any permutation group must act primitively on the minimal blocks that it admits. It is well-known that a graph Γ is isomorphic to a Cayley graph on a group G if and only if $\mathrm{Aut}(\Gamma)$ contains a regular subgroup isomorphic to G. Thus, the automorphism group of a circulant graph always acts as one of the above groups on the minimal blocks that it admits. Understanding these possible actions therefore goes a long way towards reconstructing (recursively) the full action of the automorphism group. In particular, to determine other regular subgroups, it will be important to have a good understanding of conjugacy classes in these groups. This is the goal of the next few results in this section.

Both [9, Corollary 2] or [12, Corollary 1.2] show that if $\mathrm{PGL}(d,q) \leq K \leq \mathrm{P\Gamma L}(d,q)$, then every regular cyclic subgroup of K is a Singer subgroup (and so any two are conjugate in $\mathrm{PGL}(d,q)$) unless $d = 2$ and $q = 8$, in which case $n = 9$.

The following result is well-known.

Lemma 2.2. *Two elements of* $\mathrm{Alt}(n)$ *are conjugate in* $\mathrm{Sym}(n)$ *but not in* $\mathrm{Alt}(n)$ *if and only if their cycle structures are the same, they have no cycle of even length, and the lengths of all of their odd cycles are distinct.*

Lemma 2.3. *Let* $K, K' \leq G$ *be conjugate in* G. *If* $N \triangleleft G$ *and* $KN/N = G/N$, *then there exists* $n \in N$ *with* $n^{-1}Kn = K'$.

Proof. Let $g \in G$ such that $g^{-1}Kg = K'$. As $KN/N = G/N$ there exists $k \in K$ such that $kN = gN$. Then $N = k^{-1}gN$, or equivalently, $n = k^{-1}g \in N$. Now $n^{-1}Kn = g^{-1}kKk^{-1}g = g^{-1}Kg = K'$.

Corollary 2.4. *Any two regular cyclic subgroups of* $\mathrm{PGL}(d,q)$ *are conjugate by an element of* $\mathrm{PSL}(d,q)$, *provided that* $(d,q) \neq (2,8)$. *Also, if* n *is even, then any two regular cyclic subgroups of* $\mathrm{Sym}(n)$ *are conjugate by an element of* $\mathrm{Alt}(n)$.

Proof. By [9, Corollary 2] or [12, Corollary 1.2], there is one conjugacy class of regular cyclic subgroups of $\mathrm{PGL}(d,q)$ whenever $(d,q) \neq (2,8)$, and clearly there is always one conjugacy class of regular cyclic subgroups in $\mathrm{Sym}(n)$. If K is a regular cyclic subgroup of $\mathrm{PGL}(d,q)$, then $K\mathrm{PSL}(d,q) = \mathrm{PGL}(d,q)$. Similarly, if K is a regular cyclic subgroup of $\mathrm{Sym}(n)$ where n is even, then $K\mathrm{Alt}(n) = \mathrm{Sym}(n)$. The result follows by Lemma 2.3, with $\mathrm{PSL}(d,q)$ or $\mathrm{Alt}(n)$ taking the role of N.

The concept of Ω-step imprimitivity will be important in this paper. Intuitively, on a set of cardinality n, the action of a transitive group is Ω-step imprimitive if there is a sequence of nested block systems that is as long as possible (given n). The terms "nested" and "as long as possible" may not be clear, so we provide formal definitions below, including an explicit formula for $\Omega = \Omega(n)$.

Definition 2.5. *Let G be a transitive permutation group. Let Y be the set of all block systems of G. Define a partial order on Y by $\mathcal{B} \leq \mathcal{C}$ if and only if every block of $\mathcal{C}$ is a union of blocks of $\mathcal{B}$. We say that a strictly increasing sequence of $m+1$ block systems under this partial order is an* ***m*-step imprimitivity sequence** *admitted by G.*

An m-step imprimitivity sequence is what we referred to in our intuitive description as a "nested" sequence.

Definition 2.6. *Let G be a transitive group of degree n. Let $n = \Pi_{i=1}^{r} p_i^{a_i}$ be the prime factorization of n and let $\Omega = \Omega(n) = \sum_{i=1}^{r} a_i$. Then G is* **Ω-step imprimitive** *if it admits an Ω-step imprimitivity sequence.*

A block system $\mathcal{B}$ will be said to be **normal** *if $\mathcal{B}$ is formed by the orbits of a normal subgroup. We will say that G is* **normally Ω-step imprimitive** *if G is Ω-step imprimitive with a sequence in which each block system is normal.*

Let $\mathcal{B}_0 < \ldots < \mathcal{B}_\Omega$ be an Ω-step imprimitivity sequence of G, where G is acting on X. Then $\mathcal{B}_0$ consists of singleton sets, $\mathcal{B}_\Omega = \{X\}$, and if $B_i \in \mathcal{B}_i$ and $B_{i+1} \in \mathcal{B}_{i+1}$, then $|B_{i+1}|/|B_i|$ is a prime. Thus it is not possible to have a k-step imprimitivity sequence for any $k > \Omega$, satisfying our intuitive description of the system as being "as long as possible".

Recall that we would like to characterise graphs Γ with $G, G' \leq \mathrm{Aut}(\Gamma)$, where G and G' are regular (and nonisomorphic), and G is cyclic. Our method will be to find a subgroup N of $\mathrm{Aut}(\Gamma)$ that is normally $\Omega(n)$-step imprimitive. If we can then find regular subgroups $H, H' \leq N$ with $H \cong G$ and $H' \cong G'$, then we will be able to use the fact that H and H' also admit the many block systems of N to determine a lot about the structure of Γ.

In fact, in Lemma 2.11, we will show that there is some conjugate $\delta^{-1}G'\delta \leq \mathrm{Aut}(\Gamma)$ of G', such that $K = \langle G, \delta^{-1}G'\delta\rangle$ is normally $\Omega(n)$-step imprimitive. Then in Theorem 2.14, we show that if we assume a numerical condition on n, there is a nilpotent group $N \leq K$ that contains subgroups isomorphic to both

G and G'. Clearly N is still $\Omega(n)$-step imprimitive since it must admit all of the blocks that K admits.

The next two lemmas and the definitions surrounding them are required for the proof of Lemma 2.11, which is in turn used in the proof of Theorem 2.14.

Definition 2.7. *Let G be a transitive group acting on X and $\mathcal{B}$ a block system of G. For $g \in G$ we denote by $g/\mathcal{B}$ permutation of $\mathcal{B}$ induced by g, and $G/\mathcal{B} = \{g/\mathcal{B} : g \in G\}$.*

By $\mathbf{fix}_{\boldsymbol{G}}\boldsymbol{(\mathcal{B})}$ *we mean the subgroup of G which fixes each block of $\mathcal{B}$ set-wise. That is,*

$$\mathrm{fix}_G(\mathcal{B}) = \{g \in G : g(B) = B \text{ for all } B \in \mathcal{B}\}.$$

This can also be thought of as the kernel of the projection from G to $G/\mathcal{B}$.

For $B \in \mathcal{B}$, we denote the set-wise stabilizer of the block B by $\mathbf{Stab}_{\boldsymbol{G}}\boldsymbol{(B)}$. *That is,*

$$\mathrm{Stab}_G(B) = \{g \in G : g(B) = B\}.$$

The **support of $\boldsymbol{G}$**, *denoted* $\mathrm{supp}(G)$, *is the set of all $x \in X$ that are acted on nontrivially by some $g \in G$. That is,*

$$\mathrm{supp}(G) = \{x \in X : \text{ there exists } g \in G \textit{ such that } g(x) \neq x\}.$$

Lemma 2.8. *Let $H \leq \mathrm{Sym}(n)$ admit a block system $\mathcal{B}$ and $K \triangleleft H$ such that*

1. $K \leq \mathrm{fix}_H(\mathcal{B})$,
2. *$K = T^\ell$ is a direct product of isomorphic nonabelian simple groups T, and*
3. *$K|_B \cong T$ is transitive for every $B \in \mathcal{B}$.*

Let L be a minimal nontrivial normal subgroup of K. Then $\mathrm{supp}(L)$ *is a block of H. Furthermore, if $\mathcal{C}$ is the block system of H that includes* $\mathrm{supp}(L)$, *then $\mathcal{B} \leq \mathcal{C}$.*

PROOF. Let $C = \mathrm{supp}(L)$. Note that hLh^{-1} is a minimal normal subgroup of K with $\mathrm{supp}(hLh^{-1}) = h(C)$. Furthermore, if $k \in K$ then since $L \triangleleft K$, this shows that $k(C) = C$. Hence C is a union of orbits of K, i.e. a union of blocks from $\mathcal{B}$. If we successfully show that C is a block of H, then this implies that $\mathcal{B} \leq \mathcal{C}$.

We must show that for every $h \in H$, either $C \cap h(C) = \emptyset$, or $h(C) = C$. Consider any $h \in H$ such that $C \cap h(C) \neq \emptyset$. We will show that $h(C) = C$.

If $h(C) = C$ we are finished, so towards a contradiction we assume that

$$\mathrm{supp}(hLh^{-1}) = h(C) \neq C = \mathrm{supp}(L),$$

in which case $hLh^{-1} \neq L$. Since both L and hLh^{-1} are normal in K, so is their intersection. As $K = T^\ell$, $L \cong T$ is simple; we conclude that $L \cap hLh^{-1} = 1$. The same argument shows that when we write $K = T^\ell$, the intersection of L with each copy of T is either trivial or all of T, and the same is true for hLh^{-1}. Thus, L and hLh^{-1} must commute with each other.

We have seen that for any $k \in K$, $k(C) = C$; the same argument shows that $k(h(C)) = h(C)$. Hence $k(C \cap h(C)) = C \cap h(C)$, so $C \cap h(C)$ is a union of blocks from $\mathcal{B}$. Let $B \in \mathcal{B}$ with $B \subseteq C \cap h(C)$.

Notice that $L|_B \lhd K|_B \cong T$. Since $B \subseteq C = \mathrm{supp}(L)$, this forces $L|_B = K|_B \cong T$ to be transitive on B. Furthermore this action is a faithful representation of T since the kernel of the action of $L|_B$ is a normal subgroup of the simple group T. The same is true for $hLh^{-1}|_B$. Since T is nonabelian and the actions on B are faithful and identical, L and hLh^{-1} cannot commute, a contradiction.

Lemma 2.9. *Let G, G' be regular abelian subgroups of a primitive group K of degree n, with G cyclic. Let p be any prime divisor of n, and let G'_p be any subgroup of G' of order p. Then there exists $\delta \in \mathrm{soc}(K)$ such that $G_p = \delta^{-1}G'_p\delta$.*

Proof. If $n = p$ is prime, then G and G' are Sylow p-subgroups of K and of $\mathrm{soc}(K)$, and the result follows.

If $\mathrm{PGL}(d,q) \leq K \leq \mathrm{P\Gamma L}(d,q)$ and both G and G' are Singer subgroups then by Corollary 2.4 there exists $\delta \in \mathrm{soc}(K) = \mathrm{PSL}(d,q)$ such that $\delta^{-1}G'\delta = G$, and we are done. By Theorem 2.1 the only remaining possibilities are $K = \mathrm{Alt}(n)$, $\mathrm{Sym}(n)$, or $\mathrm{PGL}(2,8) \leq K \leq \mathrm{P\Gamma L}(2,8)$.

If $\mathrm{PGL}(2,8) \leq K \leq \mathrm{P\Gamma L}(2,8)$ then $n = 9$ and both G and G' are cyclic groups of order 9. Let $G \leq P$ and $G' \leq Q$, where P and Q are Sylow 3-subgroups of $\mathrm{P\Gamma L}(2,8)$. It can be shown that any Sylow 3-subgroup of the natural action of $\mathrm{P\Gamma L}(2,8)$ on points, contains a unique semiregular subgroup of order 3. Note that in P, this subgroup is G_3 since G_3 is the unique semiregular subgroup of order 3 in G, and similarly in Q, this subgroup is G'_3. Since any conjugate of a Singer subgroup is a Singer subgroup, P and Q each contain a Singer subgroup, S and T say (respectively), and since S and T are regular of order 9, each contains a semiregular subgroup of order 3, so $G_3 \leq S$ and $G'_3 \leq T$. By Corollary 2.4, there is some $\delta \in \mathrm{PSL}(2,8)$ such that $\delta^{-1}T\delta = S$. Since the subgroup of order 3 in S is unique, we see that $G_3 = \delta^{-1}G'_3\delta$.

If $\mathrm{soc}(K) = \mathrm{Alt}(n)$, then let $G_p \leq G$, $G'_p \leq G'$ be subgroups of order p. The semiregular cyclic groups G_p and G'_p are generated by elements g and g' respectively, where $g, g' \in \mathrm{Alt}(n)$ each has a cycle structure consisting of $n/p > 1$ cycles of length p, since n is composite. By Lemma 2.2, there is some $\delta \in \mathrm{Alt}(n)$ such that $g = \delta^{-1}g'\delta$, and so $G_p = \delta^{-1}G'_p\delta$.

Let $G \leq \mathrm{Sym}(n)$ admit a block system $\mathcal{B}$. It is straightforward to observe that there is a block system $\mathcal{D}$ in $G/\mathcal{B}$ if and only if there is a block system $\mathcal{C} \succeq \mathcal{B}$ of G where a block of $\mathcal{C}$ consists of the union of all blocks of $\mathcal{B}$ contained within a block of $\mathcal{D}$.

Definition 2.10. *If G admits block systems $\mathcal{B}$ and $\mathcal{C}$ with $B \leq C$, then we denote the corresponding block system $\mathcal{D}$ in $G/\mathcal{B}$ by $\mathcal{C}/\mathcal{B}$.*

We are now ready to show that we can find a group that contains our regular cyclic subgroup and a conjugate of any other given regular abelian subgroup, and is $\Omega(n)$-step imprimitive. Although it is not immediately clear from the statements of the results as written, [18, Theorem 4.9 (i)] is a consequence of this lemma. The authors plan to explain and expand on this generalisation in a forthcoming paper.

Lemma 2.11. *Let G, G' be regular abelian subgroups of a permutation group of degree n, with G cyclic. Let $n = p_1^{a_1} \cdots p_r^{a_r}$ be the prime-power decomposition of n, and $\Omega = \Omega(n) = \sum_{i=1}^{r} a_i$. Then there exists $\delta \in \langle G, G' \rangle$ such that $\langle G, \delta^{-1}G'\delta \rangle$ is normally Ω-step imprimitive.*

Proof. We proceed by induction on $\Omega = \Omega(n)$. If $\Omega(n) = 1$, then n is prime and the result is trivial. Let G, G', and n satisfy the hypotheses, with $\Omega(n) \geq 2$. Assume the result holds for all permutation groups of degree n' with $\Omega(n')$ at most $\Omega(n) - 1$. If $H = \langle G, G' \rangle$ is primitive, then by Lemma 2.9, there exists $\delta \in H$ such that $\langle G, \delta^{-1}G'\delta \rangle$ is imprimitive, so we may assume without loss of generality that H is imprimitive.

Suppose that $\mathcal{B}$ is a block system of H with ℓ blocks of size k. We first show that if $k = p$ is prime, we can complete the proof; then we will devote the remainder of the proof to demonstrating that if k is composite, then for any prime $p \mid k$ there exists $\delta \in H$ such that $\langle G, \delta^{-1}G'\delta \rangle$ admits a normal block system with blocks of prime size p. Replacing G' by this conjugate and $\mathcal{B}$ by this system then completes the proof.

Suppose $k = p$ is prime, and set $\mathcal{B}_1 = \mathcal{B}$. By the induction hypothesis, since $\Omega(n/p) = \Omega(n) - 1 = \Omega - 1$, there exists $\delta \in H$ such that $\langle G, \delta^{-1}G'\delta \rangle/\mathcal{B}_1$ is normally $(\Omega - 1)$-step imprimitive with $(\Omega - 1)$-step imprimitivity sequence $\mathcal{B}_1/\mathcal{B}_1 < \mathcal{B}_2/\mathcal{B}_1 < \ldots < \mathcal{B}_\Omega/\mathcal{B}_1$. Then $\langle G, \delta^{-1}G'\delta \rangle$ is normally $\Omega(n)$-step imprimitive with $\mathcal{B}_0 < \mathcal{B}_1 < \ldots < \mathcal{B}_m$. This completes the proof when $k = p$.

Suppose now that k is composite. We assume that k is chosen to be minimal, and so by [2, Exercise 1.5.10] $\mathrm{Stab}_H(B)|_B$ is primitive. Since k is composite, for any block $B \in \mathcal{B}$, $\mathrm{Stab}_H(B)|_B$ is doubly-transitive as $\mathbb{Z}_k$ is a Burnside group [2, Theorem 3.5A]. Since the groups in Theorem 2.1(1) are not of composite degree, $\mathrm{Stab}_H(B)|_B$ has nonabelian simple socle, T_B. In fact, $T_B \cong \mathrm{PSL}(d, q)$ for some d, q, or $T_B \cong \mathrm{Alt}(k)$, so T_B is doubly-transitive. Since $\mathrm{fix}_H(\mathcal{B})|_B \triangleleft \mathrm{Stab}_H(B)|_B$, its intersection with T_B is nontrivial and normal in T_B; since T_B is a simple group, we conclude that $\mathrm{soc}(\mathrm{fix}_H(\mathcal{B})|_B) = T_B$. Thus $\mathrm{fix}_H(\mathcal{B})|_B$ has a doubly-transitive socle, so must itself be doubly-transitive. Let $K = \mathrm{soc}(\mathrm{fix}_H(\mathcal{B}))$. Note that $K \triangleleft H$ as K is characteristic in $\mathrm{fix}_H(\mathcal{B})$.

We claim that K, H, and $\mathcal{B}$ satisfy the hypotheses of Lemma 2.8. The parts that we have not yet shown are (2) and (3). Clearly, $K|_B \triangleleft \mathrm{fix}_H(\mathcal{B})|_B$. Thus $(K|_B \cap T_B) \triangleleft T_B$. Since T_B is simple, we either have $K|_B \geq T_B$, or $K|_B = 1$. Note that $K|_{B'}$ is conjugate in H to $K|_B$ for any $B' \in \mathcal{B}$, so the latter would imply $K = 1$, but a socle of a nontrivial finite group cannot be trivial. So $K|_B \geq T_B$. Since any socle is a direct product of simple groups, it is not possible that $K|_B > T_B$. Thus $K|_B = T_B$ (which is in fact doubly-transitive). Given that for any $B' \in \mathcal{B}$, the groups T_B and $T_{B'}$ are conjugate in H, we have hypothesis (3). The facts that any socle is a direct product of simple groups, and that $K \leq \mathrm{fix}_H(\mathcal{B})$, force hypothesis (2), although the ℓ of Lemma 2.8 may not be the same as the ℓ we have defined in this proof.

Since the hypotheses of Lemma 2.8 are satisfied, we conclude that for any minimal normal subgroup L of K, $\mathrm{supp}(L)$ is a block of H, so there exists a block system $\mathcal{C} \succeq \mathcal{B}$ with r blocks where $r \mid \ell$, such that for every $C \in \mathcal{C}$ and $t \in T$ there is $g_{t,C} \in K$ with $(g_{t,C})|_C = t$ and $(g_{t,C})|_{C'} = 1$ if $C' \neq C$. Let $\mathcal{C} = \{C_i : 1 \leq i \leq r\}$, and $\mathcal{B} = \{B_{i,j} : 1 \leq i \leq r \text{ and } 1 \leq j \leq \ell/r\}$. We additionally assume that $\cup_{j=1}^{\ell/r} B_{i,j} = C_i$. Let p be a prime divisor of k; if $T \cong \mathrm{PSL}(d,q)$ then choose p to be a divisor of $k/\gcd(d,q-1)$. Let $G_p \leq \mathrm{fix}_G(\mathcal{B})$ be the unique subgroup of G of order p, and $G'_p \leq \mathrm{fix}_{G'}(\mathcal{B})$ a subgroup of order p. Certainly, $G_p, G'_p \leq \mathrm{fix}_H(\mathcal{B})$, so $G_p|_B, G'_p|_B \leq \mathrm{fix}_H(\mathcal{B})|_B$ for any $B \in \mathcal{B}$.

For any $1 \leq i \leq r$, let $\mathrm{fix}_H(\mathcal{B})|_{B_{i,1}}$ take the role of K in Lemma 2.9. The lemma tells us that there exists $\delta_1 \in T_{B_{i,1}}$ such that $\delta_1^{-1} G'_p \delta_i|_{B_{i,1}} = G_p|_{B_{i,1}}$. Let $G'_p = \langle h \rangle$, and $G_p = \langle g \rangle$. Then there exists $b_i \in \mathbb{Z}$ such that $gh^{b_i}|_{B_{i,1}} = 1$. By the definition of C in Lemma 2.8, it must be the case that for every $1 \leq j \leq \ell/r$, $gh^{b_i}|_{B_{i,j}} = 1$, and so $\delta_i^{-1} G'_p \delta_i|_{C_i} = G_p|_{C_i}$. Then $g_{\delta_i,C_i}^{-1} G'_p g_{\delta_i,C_i}|_{C_i} = G_p|_{C_i}$, while $g_{\delta_i,C_i}^{-1} G'_p g_{\delta_i,C_i}|_{C_j} = G'_p$ for any $j \neq i$, since $g_{\delta_i,C_i}|_{C_j} = 1$. So if $\delta = \Pi_{i=1}^{r} g_{\delta_i,C_i}$ then $\delta \in K$ and $\delta^{-1} G'_p \delta = G_p$ is a central subgroup of $\langle G, \delta^{-1} G' \delta \rangle$, whose orbits are blocks of size p. Thus, replacing G' with $\delta^{-1} G' \delta$, we assume without loss of generality that $k = p$.

Definition 2.12. *Let $K \leq \mathrm{Sym}(n)$ contain a regular abelian subgroup G. We say that a nilpotent subgroup N with $G \leq N \leq K$* ***mimics every regular abelian subgroup of K****, if a regular abelian group $M \leq \mathrm{Sym}(n)$ is contained in K if and only if N contains a regular abelian subgroup isomorphic to M.*

We first give a sufficient condition for a group K to contain a nilpotent subgroup that mimics every regular abelian subgroup of K.

Lemma 2.13. *Let G be a regular cyclic subgroup of K. Suppose that whenever $R \leq K$ is a regular abelian subgroup, then there exists $\delta \in K$ such that $\langle G, \delta^{-1} R \delta \rangle$ is nilpotent. If N is a maximal nilpotent subgroup that contains G, then N mimics every regular abelian subgroup of K.*

Proof. Let N be a maximal nilpotent subgroup that contains G, and $R \leq K$ a regular abelian subgroup of K. We will show that $\delta \in K$ with $\delta^{-1} R \delta \leq N$, which will establish the result.

Let $n = p_1^{a_1} \cdots p_r^{a_r}$ be the prime-power decomposition of n. Note that since G is a regular cyclic group, for any $1 \leq i \leq r$ it admits block systems $\mathcal{B}_i$ with blocks of size $p_i^{a_i}$ and $\mathcal{C}_i$ with blocks of size $n/p_i^{a_i}$, and if a group containing G as a subgroup admits a block system whose blocks have one of these sizes, it must be one of these block systems. As N is nilpotent and $G \leq N$, by [3, Lemma 10] for every $1 \leq i \leq r$, N admits $\mathcal{B}_i$ and $\mathcal{C}_i$ as normal block systems. Similarly, $\langle G, R \rangle$ is nilpotent and contains G, so admits each $\mathcal{B}_i$ and $\mathcal{C}_i$. Then $\langle R, N \rangle = \langle G, R, N \rangle$ admits $\mathcal{B}_i$ and $\mathcal{C}_i$ as block systems, and so $\langle R, N \rangle \leq \Pi_{i=1}^{r} \mathrm{Sym}(p_i^{a_i})$.

Now, $R/\mathcal{C}_1$ and $N/\mathcal{C}_1$ are p_1-subgroups of $\mathrm{Sym}(p_1^{a_1})$, and so there exists $\omega_1 \in \langle R, N\rangle$ such that $\omega_1^{-1}R\omega/\mathcal{C}_1$ and $N/\mathcal{C}_1$ are contained in the same Sylow p_1-subgroup. Then $\langle \omega^{-1}R\omega, N\rangle/\mathcal{C}_1$ is a p_1-group. Inductively suppose that $\langle \omega_j^{-1}\dots\omega_1^{-1}R\omega_1\dots\omega_j, N\rangle/\mathcal{C}_i$ is a p_i-group for each $1 \le i \le j < r$. Then there is some $\omega_{j+1} \in \langle \omega_j^{-1}\dots\omega_1^{-1}R\omega_1\dots\omega_j, N\rangle$ such that (by conjugation of Sylow p_{j+1}-subgroups) $\langle \omega_{j+1}^{-1}\dots\omega_1^{-1}R\omega_1\dots\omega_{j+1}, N\rangle/\mathcal{C}_{j+1}$ is a p_{j+1}-group. As $\omega_{j+1} \in \langle \omega_j^{-1}\dots\omega_1^{-1}R\omega_1\dots\omega_j, N\rangle$, our inductive hypothesis implies that for each $1 \le i \le j < r$, $\langle \omega_{j+1}^{-1}\dots\omega_1^{-1}R\omega_1\dots\omega_{j+1}, N\rangle$ is still a p_i-group, completing the induction. Let $\delta = \omega_1\dots\omega_r \in K$, so that $\langle \delta^{-1}R\delta, N\rangle/\mathcal{C}_i$ is a p_i-group for every $1 \le i \le r$. This shows that $\langle \delta^{-1}R\delta, N\rangle$ is a direct product of its Sylow p-subgroups, so is nilpotent. By the maximality of N, we have $\langle \delta^{-1}R\delta, N\rangle = N$, meaning $\delta^{-1}R\delta \le N$, as required.

We now show that under a numerical condition on n, we can find a nilpotent group that has this extremely useful property.

Theorem 2.14. *Let $k = p_1\dots p_r$ be such that $\gcd(k, \varphi(k)) = 1$ where each p_i is prime, and $n = p_1^{a_1}\dots p_r^{a_r}$. Let G be a regular cyclic subgroup of a group K of degree n. Then there exists a nilpotent subgroup N, with $G \le N \le K$, that is $\Omega(n)$-step imprimitive, and mimics every regular abelian subgroup of K.*

Proof. Let $R \le K$ be a regular abelian group. By Lemma 2.11 there exists $\delta_1 \in K$ such that $\langle G, \delta_1^{-1}R\delta_1\rangle$ is normally $\Omega(n)$-step imprimitive. By [3, Theorem 12], (applying this requires our hypothesis that $\gcd(k, \varphi(k)) = 1$), there exists $\delta_2 \in \langle G, \delta_1^{-1}R\delta_1\rangle \le K$ such that $\langle G, \delta_2^{-1}\delta_1^{-1}R\delta_1\delta_2\rangle$ is nilpotent. By Lemma 2.13, any maximal nilpotent subgroup N that contains G mimics every regular abelian subgroup of K. By [3, Lemma 9], any such N is $\Omega(n)$-step imprimitive.

3. Graph Theoretic Results

Lemma 3.1. *Let p be prime and $k_1, \dots, k_j$ be positive integers. Then there exists a vertex-transitive digraph Γ whose automorphism group is $\mathbb{Z}_{p^{k_1}} \wr \dots \wr \mathbb{Z}_{p^{k_j}}$.*

Furthermore, we can choose $\Gamma = \Gamma_1 \wr \dots \wr \Gamma_j$, where each Γ_i is a circulant digraph on $\mathbb{Z}_{p^{k_i}}$.

Proof. Let $\vec{D_k}$ denote the directed cycle of length k. When $k > 2$, $\vec{D_k}$ has automorphism group $\mathbb{Z}_k \ncong \mathrm{Sym}(k)$. When $k = 2$, $\mathrm{Aut}(K_2) = \mathrm{Aut}(\bar{K_2}) = \mathbb{Z}_2 = \mathrm{Sym}(2)$.

For each $1 \le i \le j$, let

$$\Gamma_i = \begin{cases} \vec{D_{p^{k_i}}} \text{ if } p^{k_i} > 2; \\ K_2 \text{ if } p^{k_i} = 2, \text{ and either } i = 1 \text{ or } \Gamma_{i-1} \ne K_2; \text{ and} \\ \bar{K_2} \text{ otherwise.} \end{cases}$$

Let $\Gamma = \Gamma_1 \wr \dots \wr \Gamma_j$. By [6, Theorem 5.7], $\mathrm{Aut}(\Gamma) = \mathbb{Z}_{p^{k_1}} \wr \dots \wr \mathbb{Z}_{p^{k_j}}$.

Definition 3.2. *Let X be a finite set and let G be a permutation group on x. The* ***2-closure*** *of G, denoted $G^{(2)}$, is the set*

$$\{\pi \in \mathrm{Sym}(X) \mid \forall (x,x') \in X^2, \textit{there exists } g_{xx'} \in G \textit{ with } (x,x')^\pi = (x,x')^{g_{xx'}}\},$$

where $\mathrm{Sym}(X)$ *is the symmetric group on X. Observe that in the definition of $G^{(2)}$, the element $g_{xx'}$ of G may depend upon the ordered pair (x,x'). The group G is said to be 2-closed if $G = G^{(2)}$.*

Lemma 3.3. *The 2-closure of any nilpotent group N, is itself nilpotent.*

Proof. To see this, we will use the fact that a group is nilpotent if and only if it is the direct product of its Sylow subgroups. Thus, $N = P_1 \times \ldots \times P_r$.

By [1, Theorem 5.1], we have $N^{(2)} = P_1^{(2)} \times \ldots \times P_r^{(2)}$. It is well-known that the 2-closure of a p-group is a p-group (see for example [2, Exercise 2.4.5(iv)]). Thus each $P_i^{(2)}$ is a Sylow subgroup of $N^{(2)}$, so $N^{(2)}$ is nilpotent.

Definition 3.4. *The* ***orbital digraphs*** *of G are the collection of digraphs whose vertices are the elements of X. Each orbital digraph has for its arcs one orbit of the ordered pairs of elements of X, under the action of G.*

It is easy to verify that $G^{(2)}$ is a subgroup of $\mathrm{Sym}(X)$ containing G and, in fact, $G^{(2)}$ is the smallest (with respect to inclusion) subgroup of $\mathrm{Sym}(X)$ that preserves every orbital digraph of G. Equivalently, $G^{(2)}$ is the automorphism group of the Cayley colour digraph obtained by assigning a unique colour to the edges of each orbital digraph of G. It follows that the automorphism group of a graph is 2-closed.

Corollary 3.5. *Let p be prime and G, H regular abelian groups of degree p^k with G cyclic. Then there exists $H' \leq \langle G, H\rangle$ such that $H' \cong H$ is regular and $(\langle G, H'\rangle)^{(2)} \cong \mathbb{Z}_{p^{k_1}} \wr \ldots \wr \mathbb{Z}_{p^{k_m}}$ for some $k_1, \ldots, k_m$ with $k_1 + \ldots + k_m = k$.*

Proof. Let Γ be the Cayley colour digraph formed by assigning a unique colour to the edges of each orbital digraph of $\langle G, H\rangle$. Since $\mathrm{Aut}(\Gamma)$ has regular subgroups isomorphic to G and to H, we see that Γ is a Cayley colour digraph on both of these groups, similar to condition (1) of Theorem 1.4.

Although the theorem is not stated for colour digraphs, most of the proof involves only permutation groups, so it is not hard to see that the same result is true for colour digraphs. By Theorem 1.4 (3), we can find $k_1', \ldots, k_{m'}'$ such that $\Gamma \cong U_1 \wr \ldots \wr U_{m'}$ where each U_i is a circulant colour digraph on $\mathbb{Z}_{p^{k_i'}}$, and $\mathbb{Z}_{p^{k_1'}} \times \ldots \times \mathbb{Z}_{p^{k_{m'}'}} \preceq_p H$. It is a standard and straightforward observation that $(\langle G, H\rangle)^{(2)} = \mathrm{Aut}(\Gamma) \geq \mathrm{Aut}(U_1) \wr \ldots \wr \mathrm{Aut}(U_{m'}) \geq \mathbb{Z}_{p^{k_1'}} \wr \ldots \wr \mathbb{Z}_{p^{k_{m'}'}}$. Call this last group K. By Lemma 3.1, K is 2-closed.

Since $\mathbb{Z}_{p^{k_1'}} \times \ldots \times \mathbb{Z}_{p^{k_{m'}'}} \preceq_p H$, the group K contains a regular subgroup $H' \cong H$. Repeating the argument to this point, with H' taking the role of H, we obtain $k_1, \ldots, k_m$ with $K' = \mathbb{Z}_{p^{k_1}} \wr \ldots \wr \mathbb{Z}_{p^{k_m}} \leq (\langle G, H'\rangle)^{(2)}$ and $\mathbb{Z}_{p^{k_1}} \times$

$\ldots \times \mathbb{Z}_{p^{k_m}} \preceq_p H'$. Furthermore, since $G, H' \leq K$ and K is 2-closed, we have $(\langle G, H'\rangle)^{(2)} \leq K$. Thus $K' \leq K$, which implies $\mathbb{Z}_{p^{k_1}} \times \ldots \times \mathbb{Z}_{p^{k_m}} \preceq_p \mathbb{Z}_{p^{k'_1}} \times \ldots \times \mathbb{Z}_{p^{k'_{m'}}}$. This in turn means $H' \leq K'$. Now we have $G, H' \leq K'$, and K' is also 2-closed by Lemma 3.1, so $(\langle G, H'\rangle)^{(2)} \leq K'$. Hence $K' = (\langle G, H'\rangle)^{(2)}$.

Given regular groups G_1 and H_1 of degree a and G_2 and H_2 of degree b, there is an obvious method for constructing digraphs that are simultaneously Cayley digraphs of $G_1 \times G_2$ and of $H_1 \times H_2$. Namely, construct a Cayley digraph Γ_1 of order a that is a Cayley digraph of G_1 and H_1 and a digraph Γ_2 of order b that is a Cayley digraph of G_2 and H_2, and then consider some sort of "product construction" of Γ_1 and Γ_2 to produce a digraph Γ of order ab with $\mathrm{Aut}(\Gamma_1) \times \mathrm{Aut}(\Gamma_2) \leq \mathrm{Aut}(\Gamma)$. We write "product construction" as there are two obvious products of Γ_1 and Γ_2 that ensure that $\mathrm{Aut}(\Gamma_1) \times \mathrm{Aut}(\Gamma_2) \leq \mathrm{Aut}(\Gamma)$: the wreath product, and the Cartesian product.

Definition 3.6. *Let* $\Gamma_1, \ldots, \Gamma_r$ *be digraphs. We say that* Γ *is of* **product type** $\Gamma_1, \ldots, \Gamma_r$ *if* $\mathrm{Aut}(\Gamma_1) \times \ldots \times \mathrm{Aut}(\Gamma_r) \leq \mathrm{Aut}(\Gamma)$.

Theorem 3.7. *Let* $n = p_1^{a_1} \ldots p_r^{a_r}$. *Let* G, H *be regular abelian groups of degree* n *with* G *cyclic and let* G_{p_i}, H_{p_i} *be Sylow* p_i*-subgroups of* G *and* H *respectively, and* Γ *a Cayley digraph on* G. *If*

> Γ *is of product type* $\Gamma_1, \ldots, \Gamma_r$, *where each* Γ_i *is a Cayley digraph on both* G_{p_i} *and a group isomorphic to* H_{p_i}, $1 \leq i \leq r$,

then

> Γ *is isomorphic to a Cayley digraph of* H.

Furthermore, the converse holds whenever $G, H \leq N \leq \mathrm{Aut}(\Gamma)$ *for some nilpotent group* N, *and we can also conclude that each* $\mathrm{Aut}(\Gamma_i)$ *is a (possibly trivial) multiwreath product of cyclic groups.*

Proof. First suppose that Γ is of product type $\Gamma_1, \ldots, \Gamma_r$. Then $\mathrm{Aut}(\Gamma_1) \times \ldots \times \mathrm{Aut}(\Gamma_r) \leq \mathrm{Aut}(\Gamma)$. For each i, there is some $H'_{p_i} \cong H_{p_i}$ such that $H'_{p_i} \leq \mathrm{Aut}(\Gamma_i)$. Thus $H \cong H'_{p_1} \times \ldots \times H'_{p_r} \leq \mathrm{Aut}(\Gamma)$, so Γ is isomorphic to a Cayley digraph of H.

Conversely, suppose that Γ is a Cayley digraph on G that is also isomorphic to a Cayley digraph of the abelian group H. Then G and some $H' \cong H$ are regular subgroups of $\mathrm{Aut}(\Gamma)$. By assumption, $G, H' \leq N \leq \mathrm{Aut}(\Gamma)$ for some nilpotent group N. By Lemma 3.3, $N^{(2)}$ is nilpotent. We can therefore choose $N = \langle G, H'\rangle^{(2)}$.

Let $N = P_1 \times \ldots \times P_r$ where P_i is a Sylow p_i-subgroup of N. Notice that for each i, $P_i = \langle G_{p_i}, H'_{p_i}\rangle^{(2)}$ since the 2-closure of a p-group is a p-group. Consider some orbit B_i of P_i. As an orbit of a normal subgroup, B_i is a block of N. By Corollary 3.5, there exists $H''_{p_i}|_{B_i} \leq \langle G_{p_i}|_{B_i}, H'_{p_i}|_{B_i}\rangle$ such that $H''_{p_i}|_{B_i} \cong H'_{p_i}|_{B_i}$ acts regularly, and $(\langle G_{p_i}|_{B_i}, H''_{p_i}|_{B_i}\rangle)^{(2)}$ is a multiwreath product of cyclic p_i-groups.

By Lemma 3.1 there exists a vertex-transitive digraph Γ_i with $\mathrm{Aut}(\Gamma_i) = (\langle G_{p_i}|_{B_i}, H''_{p_i}|_{B_i}\rangle)^{(2)}$. Now, $(\langle G_{p_i}|_{B_i}, H''_{p_i}|_{B_i}\rangle)^{(2)} \leq (P_i|_{B_i})^{(2)} = P_i|_{B_i}$ by Wielandt's Dissection Theorem [19, Theorem 6.5], so $\mathrm{Aut}(\Gamma_i) \leq P_i|_{B_i}$. We claim that $\mathrm{Aut}(\Gamma_1) \times \ldots \times \mathrm{Aut}(\Gamma_r) \leq N \leq \mathrm{Aut}(\Gamma)$, so that Γ is of product type $\Gamma_1, \ldots, \Gamma_r$. This is because for any i, $N = P_i \times N'_i$, so we can define $\mathrm{Aut}(\Gamma_i)$ to act semiregularly by commuting with every element of N'_i, and in this form, $\mathrm{Aut}(\Gamma_i) \leq P_i$ for each i.

To complete the proof, notice for each i that since $\mathrm{Aut}(\Gamma_i)$ contains regular subgroups isomorphic to $G_{p_i}|_{B_i}$ (which is cyclic) and to $H''_{p_i}|_{B_i} \cong H'_{p_i}|_{B_i}$, the digraph Γ_i is a circulant digraph of order $p_i^{a_i}$ that is also a isomorphic to a Cayley digraph of H_{p_i}.

Although the condition requiring a nilpotent group to achieve the converse in the above theorem may seem quite limiting, we observe that in Muzychuk's solution to the isomorphism problem for circulant digraphs [17], he shows that when $G \cong H$ are cyclic groups, the isomorphism problem for general n can be reduced to the prime power cases; this implies that G and a conjugate of H always lie in some nilpotent group together.

We point out in the coming corollary that for some values of n, the nilpotent group required to achieve the converse of the above theorem will always exist, even if G and H are not both cyclic.

Corollary 3.8. *Let $k = p_1 \ldots p_r$ be such that $\gcd(k, \varphi(k)) = 1$ where each p_i is prime, and $n = p_1^{a_1} \ldots p_r^{a_r}$. Let G, H be regular abelian groups of degree n with G cyclic and let G_{p_i}, H_{p_i} be Sylow p_i-subgroups of G and H respectively, and Γ a Cayley digraph on G. Then Γ is isomorphic to a Cayley digraph of H if and only if Γ is of product type $\Gamma_1, \ldots, \Gamma_r$, where each Γ_i is a Cayley digraph on both G_{p_i} and a group isomorphic to H_{p_i}, $1 \leq i \leq r$.*

Proof. The first implication is already proven in Theorem 3.7.

Conversely, suppose that Γ is a Cayley digraph on G that is also a Cayley digraph of the abelian group H. Then G and some $H' \cong H$ are regular subgroups of $\mathrm{Aut}(\Gamma)$. By Theorem 2.14, there exists a nilpotent subgroup $N \leq \mathrm{Aut}(\Gamma)$ that contains G and a regular subgroup H' isomorphic to H. Replacing H by H' and applying Theorem 3.7 yields the result.

Definition 3.9. *Let G and H be abelian groups of order n, and for each prime $p_i|n$, denote a Sylow p_i-subgroup of G or H by G_{p_i} or H_{p_i}, respectively. Define a partial order $\preceq$ on the set of all abelian groups of order n by $G \preceq H$ if and only if $G_{p_i} \preceq_{p_i} H_{p_i}$ for every prime divisor $p_i|n$.*

Remark 3.10. *If, in Theorem 3.7, H is chosen to be minimal with respect to $\preceq$, then the rank of each Sylow p_i-subgroup of H (i.e. the number of elements in any irredundant generating set) will be equal to the number of factors in the multiwreath product $\mathrm{Aut}(\Gamma_i)$. Moreover, $\mathrm{Aut}(\Gamma_1) \times \ldots \times \mathrm{Aut}(\Gamma_r)$ will contain a regular subgroup isomorphic to the abelian R if and only if $\mathrm{Aut}(\Gamma)$ contains a regular subgroup isomorphic to the abelian group R.*

Theorem 3.11. *Let $\Gamma = \mathrm{Cay}(G,S)$ for some abelian group G of order n. Let R, R' be regular abelian subgroups of $\mathrm{Aut}(\Gamma)$, with R cyclic. Suppose that there is a nilpotent group N with $R' \leq N \leq \mathrm{Aut}(\Gamma)$, such that N mimics every regular abelian subgroup of $\mathrm{Aut}(\Gamma)$. Then the following are equivalent:*

1. *The digraph Γ is isomorphic to a Cayley digraph on both R and H, where H is a regular abelian group; furthermore, if H is chosen to be minimal with respect to the partial order $\preceq$ from amongst all regular abelian subgroups of $\mathrm{Aut}(\Gamma)$, then for each i, the Sylow p_i-subgroup of H has rank m_i.*
2. *Let P_i be a Sylow p_i-subgroup of G. There exist a chain of subgroups $P_{i,1} \leq \ldots \leq P_{i,m_i-1}$ in P_i such that*
   - (a) *$P_{i,1}, P_{i,2}/P_{i,1}, \ldots, P_i/P_{i,m_i-1}$ are cyclic p_i-groups;*
   - (b) *$P_{i,1} \times P_{i,2}/P_{i,1} \times \ldots \times P_i/P_{i,m_i-1} \preceq_p H_i$, where H_i is a Sylow p_i-subgroup of H;*
   - (c) *For all $s \in S\backslash(P_{i,j} \times G'_i)$, we have $sP_{i,j} \subseteq S$, for $j = 1, ..., m_i - 1$, where G'_i is a Hall p'_i-subgroup of G (of order $n/p_i^{a_i}$). That is, $S\backslash(P_{i,j} \times G'_i)$ is a union of cosets of $P_{i,j}$.*
3. *The digraph Γ is of product type $\Gamma_1, \ldots, \Gamma_r$, where each $\Gamma_i \cong U_{i,m_i} \wr \ldots \wr U_{i,1}$ for some Cayley digraphs $U_{i,1}, ..., U_{i,m_i}$ on cyclic p_i-groups $K_{i,1}, ..., K_{i,m_i}$ such that $K_{i,1} \times \ldots \times K_{i,m_i} \preceq_p H_i$, where H_i is a Sylow p_i-subgroup of H. These in turn imply:*
4. *Γ is isomorphic to Cayley digraphs on every abelian group of order n that is greater than H in the partial order $\preceq$.*

Proof. Throughout this proof, let $n = p_1^{a_1} \ldots p_r^{a_r}$.

(1)⇒(2): By hypothesis there is a transitive nilpotent subgroup N of $\mathrm{Aut}(\Gamma)$ that contains regular subgroups isomorphic to G, R, and H. By Theorem 3.7, there exist $\Gamma_1, \ldots, \Gamma_r$ where each Γ_i is a circulant digraph of order $p_i^{a_i}$, such that Γ is of product type $\Gamma_1, \ldots, \Gamma_r$, and each $\mathrm{Aut}(\Gamma_i)$ is a (possibly trivial) multiwreath product of cyclic groups. Additionally, by Remark 3.10, we may assume that R'' is a regular abelian subgroup of $\mathrm{Aut}(\Gamma)$ if and only if $\mathrm{Aut}(\Gamma_1) \times \ldots \times \mathrm{Aut}(\Gamma_r)$ contains a regular abelian subgroup isomorphic to R''.

Let $\mathrm{Aut}(\Gamma_i) = \mathbb{Z}_{p_i^{b_{m_i,i}}} \wr \cdots \wr \mathbb{Z}_{p_i^{b_{1,i}}}$ (by Remark 3.10, this multiwreath product does have m_i factors). Then $\mathrm{Aut}(\Gamma_i)$ admits block systems $\mathcal{D}_{i,j}$, $1 \leq j \leq m_i$ consisting of blocks of size $x_{i,j} = \Pi_{\ell=1}^{j} p_i^{b_{\ell,i}}$. Observe that $\mathrm{Aut}(\Gamma_i)/\mathcal{D}_{i,j} = \mathbb{Z}_{p_i^{b_{m_i,i}}} \wr \cdots \wr \mathbb{Z}_{p_i^{b_{j+1,i}}}$ and $\mathrm{fix}_{\mathrm{Aut}(\Gamma_i)}(\mathcal{D}_{i,j})|_D = \mathbb{Z}_{p_i^{b_{j,i}}} \wr \cdots \wr \mathbb{Z}_{p_i^{b_{1,i}}}$, $D \in \mathcal{D}_{i,j}$. Additionally, $\mathrm{fix}_{\mathrm{Aut}(\Gamma_i)}(\mathcal{D}_{i,j+1})/\mathcal{D}_{i,j}$ in its action on $D/\mathcal{D}_{i,j} \in \mathcal{D}_{i,j+1}/\mathcal{D}_{i,j}$ is cyclic of order $p_i^{b_{j+1,i}}$. Let P_i be a Sylow p_i-subgroup of G. Now, as G is a transitive abelian group, $P_{i,j} = \mathrm{fix}_G(\mathcal{D}_{i,j})$ is semiregular and transitive on $D \in \mathcal{D}_{i,j}$. As $\mathrm{fix}_{\mathrm{Aut}(\Gamma_i)}(\mathcal{D}_{i,j+1})/\mathcal{D}_{i,j}$ in its action on $D/\mathcal{D}_{i,j} \in \mathcal{D}_{i,j+1}/\mathcal{D}_{i,j}$ is cyclic of order $p_i^{b_{j+1,i}}$, we see that $P_{i,j+1}/P_{i,j}$ is cyclic of prime-power order, and for the same reason,

$$P_{i,1} \times P_{i,2}/P_{i,1} \times \ldots \times P_i/P_{i,m_i-1} \preceq_p H_i,$$

where H_i is a Sylow p_i-subgroup of H. For $1 \leq j \leq m_i - 1$, N admits a block system $\mathcal{C}_{i,j}$ consisting of blocks of size $x_{i,j} \cdot n/p_i^{a_i}$, as well as a block system $\mathcal{B}_{i,j}$ consisting of blocks of size $x_{i,j}$, and of course $\mathcal{B}_{i,j} \leq \mathcal{C}_{i,j}$. As Γ is of product type $\Gamma_1, \ldots, \Gamma_r$, and each Γ_i is a circulant graph, we see that $\mathrm{Aut}(\Gamma_i) \times \mathbb{Z}_{n/p_i^{a_i}} \leq \mathrm{Aut}(\Gamma)$; furthermore, $\mathrm{Aut}(\Gamma_i) \geq \mathbb{Z}_{p_i^{a_i}/x_{i,j}} \wr \mathbb{Z}_{x_{i,j}}$. Thus, $P_{i,j}|_C \leq \mathrm{Aut}(\Gamma)$ for every $C \in \mathcal{C}_{i,j}$. Thus we see that between blocks $C, C' \in \mathcal{C}_i$, we have either every directed edge from a block of $\mathcal{B}_{i,j}$ contained in C to a block of $\mathcal{B}_{i,j}$ contained in C' or no directed edges. As $\mathcal{C}_i$ is formed by the orbits of $P_{i,j} \times G_i'$ and $\mathcal{B}_{i,j}$ is formed by the orbits of $P_{i,j}$, (2) follows.

(2) $\Rightarrow$ (3) Let $\mathcal{B}_{i,j}$ be the block system of G formed by the orbits of $P_{i,j}$, $0 \leq j \leq m_i$, and $\mathcal{C}_{i,j}$ the block system of G formed by the orbits of $P_{i,j} \times G_i'$. As for all $s \in S \backslash (P_{i,j} \times G_i')$, we have $sP_{i,j} \subseteq S$, for $j = 1, \ldots, m_i - 1$, we have $\mathrm{fix}_G(\mathcal{B}_{i,j})|_{C_{i,j}} \leq \mathrm{Aut}(\Gamma)$ for every $C_{i,j} \in \mathcal{C}_{i,j}$. Note that $\mathrm{fix}_G(\mathcal{B}_{i,m_i}) = P_i$ and that $\langle \mathbb{Z}_{p_i^{a_i}}, \mathrm{fix}_G(\mathcal{B}_{i,j})|_{C_{i,j}} : C_{i,j} \in \mathcal{C}_{i,j} \rangle$ in its action on $B_{i,m_i} \in \mathcal{B}_{i,m_i}$ is $\mathbb{Z}_{p_i^{a_i - \ell_{i,j}}} \wr \mathbb{Z}_{p_i^{\ell_{i,j}}}$ where the orbits of $P_{i,j}$ have order $p_i^{\ell_{i,j}}$. Now let $n_{i,j+1} = \ell_{i,j+1} - \ell_{i,j}$, $0 \leq j \leq m_i - 1$. Then $Q_i = \langle \mathrm{fix}_G(\mathcal{B}_{i,j})|_{C_{i,j}} : C_{i,j} \in \mathcal{C}_{i,j}, 1 \leq j \leq m_i \rangle$ in its action on $B_{i,m_i} \in \mathcal{B}_{i,m_i}$ is $\mathbb{Z}_{p_i^{n_{i,m_i}}} \wr \mathbb{Z}_{p_i^{n_{i,m_i-1}}} \wr \ldots \wr \mathbb{Z}_{p_i^{n_{i,1}}}$, and $Q_i = (\mathbb{Z}_{p_i^{n_{i,m_i}}} \wr \mathbb{Z}_{p_i^{n_{i,m_i-1}}} \wr \ldots \wr \mathbb{Z}_{p_i^{n_{i,1}}}) \times 1_{\mathrm{Sym}(n/p_i^{a_i})}$. Let $K_{i,j} = \mathbb{Z}_{p_i^{n_{i,j}}}$ for each $1 \leq j \leq m_i$. Clearly, these are cyclic p_i-groups. Also, $P_{i,j+1}/P_{i,j} \cong K_{i,j+1}$ as abstract groups for $0 \leq j \leq m_i - 1$, so assumption (b) tells us that $K_{i,1} \times \ldots \times K_{i,m_i} \preceq_p H_i$. Notice that $Q_1 \times Q_2 \times \cdots \times Q_r \leq \mathrm{Aut}(\Gamma)$, and by Lemma 3.1 there exists a digraph Γ_i with $\mathrm{Aut}(\Gamma_i) = Q_i$. The graphs $U_{i,j}$ are given in the "Furthermore" of Lemma 3.1. Thus Γ is of product type $\Gamma_1, \ldots, \Gamma_r$, completing this part of the proof.

(3) $\Rightarrow$ (1), (4) For each i, $\mathrm{Aut}(\Gamma_i)$ is a multiwreath product of m_i Cayley digraphs on cyclic groups. It is straightforward to verify that $\mathbb{Z}_a \wr \mathbb{Z}_b$ contains regular subgroups isomorphic to both $\mathbb{Z}_{ab}$ and $\mathbb{Z}_a \times \mathbb{Z}_b$.

By assumption, $\mathrm{Aut}(\Gamma_i) \geq \mathbb{Z}_{p_i^{n_{i,m_i}}} \wr \ldots \wr \mathbb{Z}_{p_i^{n_{i,1}}}$. Hence the above facts tell us that $H_i = \mathbb{Z}_{p_i^{n_{i,1}}} \times \ldots \times \mathbb{Z}_{p_i^{n_{m_i}}} \leq \mathrm{Aut}(\Gamma_i)$, and also that $H_i' \leq \mathrm{Aut}(\Gamma_i)$ for every $H_i' \succ_p H_i$. Therefore $H = H_1 \times \ldots \times H_r \leq \mathrm{Aut}(\Gamma)$, and also $H' \leq \mathrm{Aut}(\Gamma)$ for every $H' \succ H$.

The following result is obtained from the previous result by applying Theorem 2.14.

Corollary 3.12. *Let $k = p_1 \ldots p_r$ be such that* $\gcd(k, \varphi(k)) = 1$ *where each p_i is prime, and $n = p_1^{a_1} \ldots p_r^{a_r}$. Let $\Gamma = \mathrm{Cay}(G, S)$ for some abelian group G of order n. Then the following are equivalent:*

1. *The digraph Γ is isomorphic to a Cayley digraph on both $\mathbb{Z}_n$ and H, where H is a regular abelian group; furthermore, if H is chosen to be minimal with respect to the partial order $\preceq$ from amongst all regular abelian subgroups of* $\mathrm{Aut}(\Gamma)$, *then for each i, the Sylow p_i-subgroup of H has rank m_i.*

2. *Let P_i be a Sylow p_i-subgroup of G. There exist a chain of subgroups $P_{i,1} \leq \ldots \leq P_{i,m_i-1}$ in P_i such that*
   (a) *$P_{i,1}, P_{i,2}/P_{i,1}, \ldots, P_i/P_{i,m_i-1}$ are cyclic p_i-groups;*
   (b) *$P_{i,1} \times P_{i,2}/P_{i,1} \times \ldots \times P_i/P_{i,m_i-1} \preceq_p H_i$, where H_i is a Sylow p_i-subgroup of H;*
   (c) *For all $s \in S \backslash (P_{i,j} \times G'_i)$, we have $sP_{i,j} \subseteq S$, for $j = 1, ..., m_i - 1$, where G'_i is a Hall p'_i-subgroup of G (of order $n/p_i^{a_i}$). That is, $S \backslash (P_{i,j} \times G'_i)$ is a union of cosets of $P_{i,j}$.*
3. *The digraph Γ is of product type $\Gamma_1, \ldots, \Gamma_r$, where each $\Gamma_i \cong U_{i,m_i} \wr \ldots \wr U_{i,1}$ for some Cayley digraphs $U_{i,1}, ..., U_{i,m_i}$ on cyclic p_i-groups $K_{i,1}, ..., K_{i,m_i}$ such that $K_{i,1} \times \ldots \times K_{i,m_i} \preceq_p H_i$, where H_i is a Sylow p_i-subgroup of H. These in turn imply:*
4. *Γ is isomorphic to Cayley digraphs on every abelian group of order n that is greater than H in the partial order $\preceq$.*

A point about the previous results should be emphasized. That is, it is necessary to introduce the digraphs $\Gamma_1, \ldots, \Gamma_r$ from Lemma 3.1 - one cannot simply define $\Gamma_i = \Gamma[B_i]$ (the induced subgraph on the points of B_i), where $B_i \in \mathcal{B}_i$ and $\mathcal{B}_i$ is a block system whose blocks are the orbits of some Sylow p-subgroup of the regular cyclic subgroup. Rephrased, it is possible for $\Gamma[B_i]$ to be a Cayley digraph of more than one group even when Γ is only isomorphic to a Cayley digraph of a cyclic group, and even when the condition $\gcd(k, \varphi(k)) = 1$ is met. We give an example of such a digraph in the following result.

Example 3.13. *Let p and q be distinct primes such that $\gcd(pq, \varphi(pq)) = 1$, $S = \{(kp, 0), (1, 1) : k \in \mathbb{Z}_p\}$, and $\Gamma = \mathrm{Cay}(\mathbb{Z}_{p^2} \times \mathbb{Z}_q, S)$. The only abelian group H of order p^2q for which Γ is isomorphic to a Cayley digraph of H is the cyclic group of order p^2q. Nonetheless, let $\mathcal{B}$ be the block system of the left regular representation of $\mathbb{Z}_{p^2} \times \mathbb{Z}_q$, that has blocks of size p^2. Then for every $B \in \mathcal{B}$, the induced subdigraph $\Gamma[B]$ is a Cayley digraph on $\mathbb{Z}_{p^2}$ and on $\mathbb{Z}_p^2$, and is isomorphic to the wreath product of two circulant digraphs of order p.*

Proof. Towards a contradiction, suppose that Γ is isomorphic to a Cayley digraph of the abelian group H', where H' is not cyclic. Let G be the left regular representation of $\mathbb{Z}_{p^2} \times \mathbb{Z}_q$. Then $H' = \mathbb{Z}_p^2 \times \mathbb{Z}_q$, and by Theorem 2.14 there exists $H \cong H'$ such that $N = \langle G, H \rangle$ is nilpotent. Then N admits $\mathcal{B}$ as a block system as well as block systems $\mathcal{B}_p$ and $\mathcal{B}_q$ consisting of blocks of size p and blocks of size q, respectively. As $H \cong \mathbb{Z}_p^2 \times \mathbb{Z}_q$, $N/\mathcal{B}_q$ is a p-group that contains regular subgroups isomorphic to $\mathbb{Z}_p^2$ and $\mathbb{Z}_{p^2}$. By [8, Lemma 4] we have that $N/\mathcal{B}_q \cong \mathbb{Z}_p \wr \mathbb{Z}_p$.

For $0 \leq i \leq p^2 - 1$, let $B_{i,q} \in \mathcal{B}_q$ denote the block that consists of $\{(i, j) : j \in \mathbb{Z}_q\}$. Each vertex of $B_{0,q}$ is at the start of a unique directed path in Γ (that does not include digons) of length q (travelling by arcs that come from $(1, 1) \in S$), and each of these paths ends at a vertex of $B_{q,q}$. Thus any automorphism of Γ that fixes $B_{0,q}$ must also fix $B_{q,q}$, contradicting $K/\mathcal{B}_q \cong \mathbb{Z}_p \wr \mathbb{Z}_p$.

Finally, $\Gamma[B]$ is isomorphic to $\mathrm{Cay}(\mathbb{Z}_{p^2}, \{kp : k \in \mathbb{Z}_p^*\}) \cong \bar{K}_p \wr K_p$, so by Theorem 1.4, $\Gamma[B]$ is also a Cayley graph on $\mathbb{Z}_p^2$.

The converse though is true. That is, if the condition $\gcd(k, \varphi(k)) = 1$ is met, and Γ is a Cayley digraph of two abelian groups G and H with nonisomorphic Sylow p-subgroups G_p and H_p, respectively, and G is cyclic, then it must be the case that $\Gamma[B]$ is a wreath product, where $\mathcal{B}$ is formed by the orbits of G_p and $B \in \mathcal{B}$.

Corollary 3.14. *Let* $k = p_1 \dots p_r$ *be such that* $\gcd(k, \varphi(k)) = 1$ *where each* p_i *is prime, and* $n = p_1^{a_1} \dots p_r^{a_r}$*. Let* Γ *be a circulant graph of order* n*, and let* $\mathcal{B}_i$ *be the block system of the left regular representation of* $\mathbb{Z}_n$ *consisting of blocks of size* $p_i^{a_i}$*. If* $\Gamma[B_i]$*,* $B_i \in \mathcal{B}_i$*, is not a nontrivial wreath product and* H *is an abelian group of order* n *such that* Γ *is isomorphic to a Cayley digraph of* H*, then a Sylow* p_i*-subgroup of* H *is cyclic. Consequently, if* $\Gamma[B_i]$*,* $B_i \in \mathcal{B}_i$*, is not isomorphic to a nontrivial wreath product for any* $1 \leq i \leq r$ *then* Γ *is not isomorphic to a Cayley digraph of any noncyclic abelian group.*

PROOF. By Theorem 2.14, there is a transitive nilpotent subgroup N of $\mathrm{Aut}(\Gamma)$ that contains the left regular representation of $\mathbb{Z}_n$ as well as a regular subgroup isomorphic to H. The system $\mathcal{B}_i$ is a block system of N also, since N is nilpotent. Thus $\mathcal{B}_i$ is a block system of H. Let H_i denote a Sylow p_i-subgroup of H, and G_i a Sylow p_i-subgroup of $\mathbb{Z}_n$. If H_i is not cyclic, then the restrictions of H_i and G_i to any $B_i \in \mathcal{B}_i$ are nonisomorphic regular p-groups, so by Theorem 1.4, $\Gamma[B_i]$ is a nontrivial wreath product.

4. Future Work

The work in this paper provides a "template" that one can use to approach the problem of when a digraph is a Cayley digraph of two nonisomorphic nilpotent groups, as follows.

Let R, R' be two regular nilpotent groups of order n. If there exists $\delta \in \langle R, R' \rangle$ such that $\langle R, \delta^{-1} R' \delta \rangle$ is nilpotent, then $\langle R, \delta^{-1} R' \delta \rangle^{(2)}$ is also nilpotent, and writing $\langle R, \delta^{-1} R' \delta \rangle^{(2)}$ as $\Pi_{i=1}^r P_i$, where $P_1, \dots, P_r$ are all Sylow subgroups of $\langle R, \delta^{-1} R' \delta \rangle^{(2)}$, then each P_i is 2-closed. Furthermore, if R_{p_i} is the Sylow p_i-subgroup of R, R'_{p_i} is the Sylow p_i-subgroup of R', and P_i is the Sylow p_i-subgroup of $\langle R, \delta^{-1} R' \delta \rangle^{(2)}$, then $P_i = \langle R_{p_i}, \delta^{-1} R'_{p_i} \delta \rangle^{(2)}$. Thus, the subgraph induced on each orbit of P_i is a Cayley graph on the Sylow p_i-subgroups of both R and R'. So from a group theoretic point of view, this "reduces" the group theoretic characterization to the corresponding prime-power cases.

Of course, we would ideally like conditions on the connection set of a Cayley digraph of one group to be a Cayley digraph of another group, but at this time such conditions are only known in the prime-power case when one of the groups is cyclic. We also suspect that the conditions given in Theorem 3.11 and Corollary 3.12 only hold when one of the groups is a cyclic group, and different conditions will be needed for different choices of nilpotent or even abelian groups. So we have the following problem:

Problem 4.1. *Given p-groups P and P', determine necessary and sufficient conditions on $S \subseteq P$ so that* $\mathrm{Cay}(P, S)$ *is also isomorphic to a Cayley digraph of P'.*

So how can one determine if a δ that conjugates R' to lie in a nilpotent group with R exists? We have seen in this proof that sometimes in order to prove that they lie together in a nilpotent group, it is sufficient to show that they lie together in a group that is normally $\Omega(n)$-step imprimitive. Naively following the structure of the proof in this paper, we have the following problem:

Problem 4.2. *Determine for which regular nilpotent groups N and N' of order n there exists $\delta \in \langle N, N'\rangle$ such that $\langle N, \delta^{-1}N'\delta\rangle$ is (normally) $\Omega(n)$-step imprimitive.*

This condition is certainly a necessary condition for a nilpotent subgroup of $\langle N, N'\rangle$ to contain N and a conjugate of N', but this condition is also sufficient under the arithmetic condition in Theorem 3.12 by [5, Corollary 15]. The solution of Problem 4.2 will likely depend, like the proof of Theorem 2.1 and consequently Theorem 2.14, on the Classification of the Finite Simple Groups. In particular, it seems likely we will need at least a list of primitive groups which contain a regular nilpotent subgroup, as well as a perhaps a list of such nilpotent subgroups. It is worthwhile to point out that Liebeck, Praeger, and Saxl have determined all primitive almost simple groups which contain a regular subgroup [13, Theorem 1.1].

Finally, it is natural to ask whether the arithmetic conditions in Theorem 2.14 are in fact necessary. The answer, while not completely known, seems to depend upon the combinatorial object under consideration. In [7] examples are given of abelian groups G whose order does not satisfy this condition, that are not CI-groups with respect to ternary relational structures. The proof essentially boils down to showing that for these groups there is no solution to Problem 4.2 for $N = G$ and N' a certain conjugate of G, with δ in the larger group $\langle N, N'\rangle^{(3)}$ (note that $\langle N, N'\rangle \leq \langle N, N'\rangle^{(3)} \leq \langle N, N'\rangle^{(2)}$, and see [7] for notation and terms not defined here). However, for digraphs and graphs on these same groups, there is $\delta \in \langle N, N'\rangle^{(2)}$ as in Problem 4.2. So it seems that the arithmetic condition given in Theorem 2.14 (or something similar) are necessary from a group theoretic point of view but perhaps not in a combinatorial setting and depending on the combinatorial object under consideration.

References

[1] Peter J. Cameron, Michael Giudici, Gareth A. Jones, William M. Kantor, Mikhail H. Klin, Dragan Marušič, and Lewis A. Nowitz, *Transitive permutation groups without semiregular subgroups*, J. London Math. Soc. (2) **66** (2002), no. 2, 325–333. MR MR1920405 (2003f:20001)

[2] John D. Dixon and Brian Mortimer, *Permutation groups*, Graduate Texts in Mathematics, vol. 163, Springer-Verlag, New York, 1996. MR MR1409812 (98m:20003)

[3] Edward Dobson, *On isomorphisms of abelian Cayley objects of certain orders*, Discrete Math. **266** (2003), no. 1-3, 203–215, The 18th British Combinatorial Conference (Brighton, 2001). MR MR1991717 (2004m:20006)

[4] ______, *Automorphism groups of metacirculant graphs of order a product of two distinct primes*, Combin. Probab. Comput. **15** (2006), no. 1-2, 105–130. MR MR2195578 (2006m:05108)

[5] ______, *On the Cayley isomorphism problem for Cayley objects of nilpotent groups of some orders*, Electron. J. Combin. **21** (2014), no. 3, Paper 3.8, 15. MR 3262245

[6] Edward Dobson and Joy Morris, *Automorphism groups of wreath product digraphs*, Electron. J. Combin. **16** (2009), no. 1, Research Paper 17, 30. MR MR2475540

[7] Edward Dobson and Pablo Spiga, *CI-groups with respect to ternary relational structures: new examples*, Ars Math. Contemp. **6** (2013), no. 2, 351–364. MR 3015643

[8] Edward Dobson and Dave Witte, *Transitive permutation groups of prime-squared degree*, J. Algebraic Combin. **16** (2002), no. 1, 43–69. MR MR1941984 (2004c:20007)

[9] Gareth A. Jones, *Cyclic regular subgroups of primitive permutation groups*, J. Group Theory **5** (2002), no. 4, 403–407. MR MR1931365 (2003h:20004)

[10] Anne Joseph, *The isomorphism problem for Cayley digraphs on groups of prime-squared order*, Discrete Math. **141** (1995), no. 1-3, 173–183. MR 1336683 (96e:05071)

[11] István Kovács and Mary Servatius, *On Cayley digraphs on nonisomorphic 2-groups*, J. Graph Theory **70** (2012), no. 4, 435–448. MR 2957057

[12] Cai Heng Li, *The finite primitive permutation groups containing an abelian regular subgroup*, Proc. London Math. Soc. (3) **87** (2003), no. 3, 725–747. MR MR2005881 (2004i:20003)

[13] Martin W. Liebeck, Cheryl E. Praeger, and Jan Saxl, *Regular subgroups of primitive permutation groups*, Mem. Amer. Math. Soc. **203** (2010), no. 952, vi+74. MR 2588738

[14] Dragan Marušič and Joy Morris, *Normal circulant graphs with noncyclic regular subgroups*, J. Graph Theory **50** (2005), no. 1, 13–24. MR 2157535 (2006c:05073)

[15] Joy Morris, *Isomorphic Cayley graphs on different groups*, Proceedings of the Twenty-seventh Southeastern International Conference on Combinatorics, Graph Theory and Computing (Baton Rouge, LA, 1996), vol. 121, 1996, pp. 93–96. MR MR1431979 (97k:05102)

[16] ______, *Isomorphic Cayley graphs on nonisomorphic groups*, J. Graph Theory **31** (1999), no. 4, 345–362. MR MR1698752 (2000e:05085)

[17] M. Muzychuk, *A solution of the isomorphism problem for circulant graphs*, Proc. London Math. Soc. (3) **88** (2004), no. 1, 1–41. MR MR2018956 (2004h:05084)

[18] Mikhail Muzychuk, *On the isomorphism problem for cyclic combinatorial objects*, Discrete Math. **197/198** (1999), 589–606, 16th British Combinatorial Conference (London, 1997). MR MR1674890 (2000e:05165)

[19] H. Wielandt, *Permutation groups through invariant relations and invariant functions*, lectures given at The Ohio State University, Columbus, Ohio, 1969.